

Aberystwyth University

Response of the rumen archaeal and bacterial populations to anti-methanogenic organosulphur compounds in continuous-culture fermenters

Martínez-Fernández, Gonzalo; Abecia, Leticia; Martín-García, A. Ignacio; Ramos-Morales, Eva; Denman, Stuart E; Newbold, Charles J; Molina-Alcaide, Eduarda; Yáñez-Ruiz, David Rafael

Published in:
FEMS Microbiology Ecology

DOI:
[10.1093/femsec/fiv079](https://doi.org/10.1093/femsec/fiv079)

Publication date:
2015

Citation for published version (APA):
Martínez-Fernández, G., Abecia, L., Martín-García, A. I., Ramos-Morales, E., Denman, S. E., Newbold, C. J., Molina-Alcaide, E., & Yáñez-Ruiz, D. R. (2015). Response of the rumen archaeal and bacterial populations to anti-methanogenic organosulphur compounds in continuous-culture fermenters. *FEMS Microbiology Ecology*, 91(8), [fiv079]. <https://doi.org/10.1093/femsec/fiv079>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

**Response of the rumen archaeal and bacterial populations to anti-methanogenic
organosulphur compounds in continuous-culture fermenters**

Gonzalo Martínez-Fernández^{1,2}, Leticia Abecia¹, A. Ignacio Martín-García¹, Eva Ramos-
Morales¹, Stuart E. Denman², Charles J. Newbold³, Eduarda Molina-Alcaide¹ & David R.
Yáñez-Ruiz¹

¹Animal Nutrition, Estación Experimental del Zaidín (CSIC). C/ Camino del Jueves s/n,
Armillá, 18100, Granada, Spain.

²CSIRO, Agriculture Flagship, Queensland Bioscience Precinct, 306 Carmody Road, St.
Lucia, QLD, Australia.

³IBERS, Aberystwyth University, Aberystwyth, UK

Running Head: Anti-methanogenic compounds and rumen ecosystem

Correspondence: David R. Yáñez-Ruiz, Animal Nutrition, Estación Experimental del Zaidín
(CSIC), C/Camino del Jueves sn, 18100, Armilla, Granada, Spain. Tel.: +34 958 18 16 00;
fax: +34 958 18 16 09; e-mail: david.yanez@eez.csic.es

26 **Abstract**

27 The study of methanogenesis inhibitors effectiveness in the rumen have shown inconsistent
28 results, mainly due to poorly understood effects on the key microbial groups involved in
29 methane CH₄ synthesis pathways. This experiment was designed to assess the effect of
30 propyl propane thiosulfinate (PTS), diallyl disulfide (DDS) and bromochloromethane (BCM)
31 on rumen fermentation, methane production and microbial populations on continuous culture
32 fermenters. No effects on total VFA were observed with PTS and DDS, while were decreased
33 with BCM. Amylase activity increased with BCM compared to the other treatments. A
34 decrease on methane production was observed with PTS (48%) and BCM (94%) compared to
35 control. Concentration of methanogenic archaea decreased with BCM from day 4 onward and
36 with PTS on days 4 and 8. The pyrosequencing analysis revealed that PTS and BCM
37 decreased the relative abundance of Methanomicrobiales and increased that of
38 *Methanobrevibacter* and *Methanosphaera*. The concentration of total bacteria was not
39 modified by any treatment, although BCM treatment increased the relative abundance of
40 *Prevotella* and decreased that of *Ruminococcus*. These results suggest that the inhibition of
41 methane production in the rumen by PTS and BCM is associated with a shift in the archaeal
42 biodiversity and changes in bacterial community with BCM.

43 **Keywords:** Garlic compounds; methanogens; microbial community; rumen.

44

44 **Introduction**

45 Enteric microbial fermentation in ruminants is an important source of anthropogenic methane
46 (CH₄), a potent greenhouse gas and its production represents a loss of around 2-12% energy
47 for the animal, and consequently, a decrease of productivity (Morgavi *et al.*, 2010). The
48 microbial populations responsible for fermentation in the rumen are comprised of an
49 extremely diverse and complex mix of bacteria, protozoa, fungi and archaea. This enables the
50 animal to digest and metabolize plant structural carbohydrates that otherwise could not be
51 achieved with only its digestive enzymes. Methanogenic archaea fill the role in the rumen of
52 terminal reducers of carbon, producing CH₄ mainly from H₂ and CO₂.

53 In the last decade, a wide range of compounds has been tested for their ability to reduce
54 methane emissions (Benchaar & Greathead, 2011). Some plant secondary metabolites have
55 shown promise due to their antimicrobial activity, including garlic-derived compounds (Hart
56 *et al.*, 2008). However, inconsistent results together with adverse effects on fiber digestion
57 and fermentation through inhibition of some bacterial groups have also been reported, with
58 the magnitude of these adverse effects varying depending upon the types and doses and diet
59 composition. Part of the inconsistency in the effects has been associated with the variety of
60 compounds included in the plant extracts, which highlights the importance of using pure
61 active compounds to understand the effects. In some cases, the effect is reversed after a few
62 days of treatment (Soliva *et al.*, 2011). Thus, the research aiming to decrease CH₄ emissions
63 from ruminants has to be built upon a correct understanding on the mechanisms of action
64 involved in relation to the main microbial groups likely to be affected (bacteria and archaea).
65 We have recently observed that two organosulphur compounds, propyl propane thiosulfinate
66 (PTS) and diallyl disulphide (DDS), strongly inhibit methane production (up to 96 and 62 %,
67 respectively) in batch culture after 24 h incubation of goats' rumen fluid (Martínez-
68 Fernández *et al.*, 2013). However, the persistency of such effect needs to be confirmed and

the microbial groups involved identified. This would enable an understanding into the mechanisms of action and facilitate practical implementation as a feeding strategy. Therefore, the aim of this work was to evaluate the effects of DDS and PTS on rumen fermentation, microbial abundances and community structure and on methane production in continuous-culture fermenters (CCF) inoculated with goats' rumen fluid.

Materials and methods

Fermenters, treatments, diet and animals

Eight CCF were used following the model of Muetzel *et al.* (2009) with an effective volume of 1000 mL. The treatments were control (without additive), DDS (purity of 80%), PTS (purity of 75%) and BCM that was included as antimethanogenic reference compound (positive inhibition control). BCM was entrapped in an α -cyclodextrin matrix (May *et al.*, 1995) before being included in the diet to ensure its stability. The formulation was prepared in our laboratory as dry white powder in 1 to 2 kg batches and contained 10% to 12% (wt/wt) of BCM. The additives and doses were selected from previous results obtained in batch cultures (Martínez-Fernández *et al.*, 2013): 80 $\mu\text{L L}^{-1}$ per day for DDS, 200 $\mu\text{L L}^{-1}$ per day for PTS and 160 mg L^{-1} per day for BCM. The DDS and BCM were provided by Sigma-Aldrich Chemical (catalog numbers 317691 and 48067, respectively); PTS was provided by DMC Research Center SL (Granada, Spain). The experimental diet (Table 1) was composed of alfalfa hay and concentrate in a 50:50 ratio.

Eight Murciano-granadina goats fitted with permanent rumen cannula were used as donors of rumen content for the experiment. Goats were adapted for 21 days to the experimental diet and were fed once a day (9:00 h) with free access to water and mineral salt block (Pacsa Sanders, Sevilla, Spain). Animals were cared by trained personnel in accordance with the Spanish guidelines for animal protection (Royal decree, 2005) and the European

Convention for the Protection of Vertebrates used for Experimental and other Scientific Purposes (European Directive, 2007). All the experimental procedures involved in this study were approved (Proc. CB-INAN 2012001) by the Animal Welfare Committee at the Institute of Animal Nutrition (CSIC, Spain).

Experimental procedure and sampling

Eight CCF (1L), as described by Muetzel *et al.* (2009), were used in two replicated incubation runs of 12 days each. Eight adult Murciano-Granadina goats fitted with ruminal canula were used as donors of ruminal contents. For each incubation run, two groups of four fermenters were inoculated (700 mL per fermenter) with a different pool, each one obtained from three different animals selected randomly. This resulted in four different pools as follows: pool 1 (goats 1, 2 and 3) and pool 2 (goats 4, 5 and 6), used in the first run; pool 3 (goats 2, 5 and 7) and pool 4 (goats 3, 6 and 8), used in the second run. Treatments (control, DDS, PTS and BCM) were randomly supplied to one of the 4 fermenters inoculated with each rumen content pool, receiving each pool all the treatments. Each fermenter was fed 16 g of fresh matter per day of the basal diet ground at 1 mm, in two equal portions at 09:00 and 14:00 h. Flow through fermenters was maintained by continuous infusion of artificial saliva (Muetzel *et al.*, 2009) at a rate of 40 mL h⁻¹ and CO₂ was continuously infused to keep anaerobic conditions. Fermenters were maintained in a water bath at 39°C. On days 0, 4, 8 and 12 of incubation 10 mL of the fermenters content were collected before the morning feeding for VFA analysis and DNA extraction and kept at -20°C and -80 °C, respectively. On days 4 and 12 of incubation 1 mL of the fermenters content was collected before feeding and frozen at -80°C for determination of amylase and xylanase activities.

On day 12 of each incubation run in CCF a batch culture trial (Theodorou *et al.*, 1994), was carried out to incubate fermenters content for 24 h to measure CH₄ production. The content

of each fermenter was filtered through two layers of cheesecloth while bubbling with CO₂. The substrate incubated consisted of 500 mg of the diet fed to fermenters in 120 mL serum bottles with 60 mL of the fermenter content. Three replicates and a blank of each fermenter and treatment were used. Bottles were sealed with rubber stoppers and aluminum caps and incubated at 39°C in a water bath. At 24 h after inoculation, the total gas volume was measured in each bottle and a sample of the gas was collected in a graduated syringe and transferred to a 5 mL vacuum tube (Venoject, Terumo Europe N.V., Leuven, Belgium) and then kept at room temperature before methane concentration was measured by gas chromatography (GC).

Chemical analysis and calculations

Dry matter (DM, method ID 934.01), ash (method ID 942.05), ether extract (method ID 7.045) and crude protein (CP, method ID 984.13) in samples were determined by the procedures of the Association of Official Analytical Chemists (AOAC, 2005). Gross Energy (GE) was measured with an adiabatic calorimeter (Model 1356, Parr Instrument Co., Moline, IL). The neutral detergent fibre (NDF), acid detergent fibre (ADF) and acid detergent lignin (ADL) contents were analysed following the methodology described by Van Soest *et al.* (1991) using an ANKOM Model 220 Fiber Analyzer (Macedon, NY). The α -amylase enzyme was used for NDF analysis in the concentrate, and both NDF and ADF were expressed exclusive of residual ash. The ADL content was determined by solubilization of cellulose in the ADF residue with 72% sulphuric acid.

To measure the enzymatic activities in fermenters content, cells were lysed using a Mini-Beadbeater (BioSpec Products, Inc., Bartlesville, OK, USA) for one minute to release intracellular enzymes, during all the process the samples were kept in ice. Cell material was removed by centrifugation (10,000×g, 10 min, 4 °C) and the supernatant was used for

analyses. Xylanase (EC 3.2.1.8.) and amylase (EC 3.2.1.1.) activities were determined (Giraldo *et al.*, 2008) using oat beachwood xylan and soluble starch, respectively, as substrates. Enzymatic activities were expressed as micromoles of glucose or xylose released in 1 min from the corresponding substrates per mL of sample at 39 °C and pH 6.5.

The individual VFA concentrations were analysed using the gas chromatography technique described by Isac *et al.* (1994).

The CH₄ concentration was determined by GC using a HP Hewlett 5890, Packard Series II gas chromatograph (Waldbronn, Germany). A sample of 0.5 mL of gas was injected using a 1 mL Sample-Lock® syringe (Hamilton, Nevada, USA).

The amount of methane produced (micromoles) was calculated by multiplying the total gas produced (micromoles) with the concentration of methane obtained.

Real-Time PCR Analysis

Samples from the fermenters content were collected on days 0, 4, 8 and 12 for DNA isolation. Samples were freeze-dried and thoroughly mixed by physical disruption using a bead beater (Mini-bead beater 8, BioSpec Products, Bartlesville, USA) for 1 min before using QIAGEN QIAamp® DNA stool mini kit (Qiagen Ltd., UK) following the manufacturer's instructions but with higher temperature (95°C) for lysis incubation. The DNA samples were used as templates for quantify the copy numbers of 16S rRNA (for bacteria), methyl coenzyme M reductase A (mcrA) gene (for methanogenic archaea), and 18S rRNA (for protozoa) by real-time quantitative PCR (qPCR).

The yield and purity of the extracted DNA were assessed using NanoDrop ND-1000 Spectrophotometer (NanoDrop Technologies, Wilmington, DE).

Primer sets used were as follows: forward: 5'-GTG-STGCAYGGYTGTCTGTC-3' and reverse: 5'-ACGT- CRTCCMCACCTTCCTC-3' for total bacteria (Maeda *et al.*, 2003) and

forward: 5'-GCTTTCGWTGGTAGT-GTATT-3' and reverse: 5'-CTTGCCCTCYAATCGT-WCT-3' for protozoa (Sylvester *et al.*, 2004). The primer sets for detection and enumeration of methanogenic archaea (*mcrA*) were forward: 5'-TTCGGTGGATCD-CARAGRGC-3' and reverse: 5'-GBARGTCGWAWC- CGTAGAATCC-3' (Denman *et al.*, 2007). Three replicates of each extract were used and a negative control was loaded on each plate run to screen for possible contamination or dimer formation and to set the background fluorescence for plate normalization.

Real-time PCR analyses were performed on iQ5 multicolor Real-Time PCR Detection System (BioRad Laboratories Inc., Hercules, CA). One microliter of DNA extract was added to amplification reactions (25 μ L) containing 0.2 μ L of each primer (10 μ M) and 12.5 μ L of iQ SYBR Green Supermix (BioRad Laboratories Inc.). Cycling conditions were 95°C for 5 min; 40 cycles of 95°C for 15 s, 60°C for 30s, and 72°C for 55 s; and 72°C for 1 min. The threshold cycle (amplification cycle in which product formation exceeds background fluorescence) of each sample was determined during the exponential phase of amplification. The absolute amount for each microbial group, expressed as the number of DNA copies/g of fresh matter, was determined using standards. The qPCR standards consisted of the plasmid pCR 4-TOPO (Invitrogen, Carlsbad, CA) with an inserted 16S, *mcrA*, or 18S gene fragment corresponding to a conserved sequence of total bacteria, methanogenic archaea, or protozoa, respectively. The number of gene copies present in the plasmid extracts was calculated using the plasmid DNA concentration and the molecular mass of the vector with the insert. The concentrated plasmid was serially diluted (10-fold) to generate a standard curve.

Pyrosequencing and sequence analysis

The yield and purity of the extracted DNA from day 12 before feeding, were assessed using NanoDrop® ND-1000 Spectrophotometer (NanoDrop Technologies, Wilmington, USA). For

sequencing of the 16S rDNA gene using the Roche /454 system, previously published primers were modified to include the 454 adaptor sequences and sample specific barcodes allowing samples to be multiplexed. Amplification of the bacterial V1-V2 regions of 16S rRNA was performed using the primer pair 27F and 357R (Liu *et al.*, 2007) . The archaeal hypervariable V6 region of the 16S rRNA gene was amplified using the primer pair 958F and 1048 Rmajor (Galand *et al.*, 2009). The PCR was performed in triplicate, in a total volume of 25 μ L containing 10x PCR buffer, 10 mM dNTP mix, 10 μ M of forward and reverse primers, 1U FastStart Polymerase, and 1 μ L of DNA template. The amplification conditions were: an initial denaturation step at 95°C for 2 min; 30 cycles of denaturation at 95°C for 30s, annealing at 55°C for 30s, and elongation at 72°C for 2 min; and a final extension step at 72°C for 7 min. The size of the PCR products was then checked on a 1% agarose gel electrophoresis. Then triplicates were pooled together and products were then purified using the short fragment removal method described by Roche using their GS FLX amplicon DNA preparation guide and AMPure beads. The purified PCR products were quantified using Quant-iT PicoGreen dsDNA quantification kit (Invitrogen) and mixed in equimolar amounts to 10^7 molecules μ L⁻¹ sample. The amplicon pooled libraries were pyrosequenced on a Roche 454 FLX Titanium. The flowgram (sff) files were converted to fasta DNA (fna) and quality score (qual) file on the 454 cluster and transferred onto a Linux based workstation running the Quantitative Insights Into Microbial Ecology (QIIME) software (Caporaso *et al.*, 2010b). Sequences were filtered to exclude those with mismatches in the primer sequence, exceeding 6 homopolymer base runs or sequences containing ambiguous bases. The libraries were split according to the 10nt barcode incorporated into the forward primer. The error-corrections of amplicon pyrosequences were made using Acacia (Bragg *et al.*, 2012). The OTUs were generated by clustering at 97% sequence identity using UCLUST (Edgar, 2010) for bacteria and CD-hit for archaea (Li & Godzik, 2006). Representative sequences were aligned to the

reads of the GreenGenes database (DeSantis Jr *et al.*, 2006) using PyNAST (Caporaso *et al.*, 2010a). Taxonomic classification was assigned using the Basic Local Alignment Search Tool (BLAST). Beta diversity was used to create principal coordinate analysis (PCoA) plots using weighted UniFrac distances. Unifrac (Lozupone & Knight, 2005).

Statistical analysis

Fermentation parameters and microbial population were analysed as a repeated measures univariate analysis using GLM procedure of SPSS (IBM SPSS Statistics v.19, IBM Corp., Somers, NY). The linear model used for each dependent variable accounted for the effects of treatment (T), day (d) and Txd interaction. Effects were considered significant at $P \leq 0.05$. When significant differences were detected, differences among means were studied using the LSD comparison test. An ANOVA analysis was used to establish differences in OTUS due to the treatments using R software.

Results

Ruminal fermentation

Methane produced in batch cultures after 24 h of incubation (Table 2) was decreased ($P \leq 0.005$) by PTS (48 %) and BCM (94 %) compared with control. The pH values in fermenters were not affected ($P = 0.308$) by the treatment, although tended to decrease as incubation time progressed ($P = 0.072$). Within treatment total VFA and individual VFA molar proportions differed ($P \leq 0.044$) for days 0 and 4 and then remained unchanged from the day 4 onward. Total VFA concentration decreased ($P < 0.001$) only with BCM addition from day 4 onward compared with control. Acetate molar proportion decreased ($P \leq 0.038$) from day 4 with all the treatments compared with control. Propionate molar proportion increased ($P \leq 0.044$) with all the treatments on day 4, although from day 4 remained higher ($P < 0.001$)

only for BCM treatment. *Iso*-butyrate molar proportion was not affected ($P = 0.222$) by any of the treatments. Molar proportion of butyrate was higher ($P \leq 0.034$) with BCM compared with control and the other two treatments from day 4 onward. Valerate and *Iso*-valerate molar proportions were higher ($P < 0.001$) from day 4 and day 8 onward, respectively, with BCM compared with control and all the treatments. Acetate to propionate ratio decreased ($P < 0.001$) from day 4 onward with BCM compared with other treatments and control. Only on day 4 the ratio was higher ($P = 0.036$) for PTS and on days 4 and 12 it was lower ($P \leq 0.040$) for DDS compared to the control.

The xylanase and amylase activities (Table 3) did not change from day 4 to 12 of incubation ($P > 0.050$). Xylanase activity tended to be affected by PTS treatment ($P = 0.051$) as compared to the control and BCM and DDS. Amylase activity increased ($P < 0.022$) on days 4 and 12 with BCM compared to the control and BCM and DDS.

Microbial community

The effects of the studied additives during 12 days of treatment on the numbers of bacteria, archaea and protozoa are shown in table 4. The abundance of the total bacterial population increased ($P \leq 0.040$) for all treatments on days 4 and 12, with a further increase by BCM treatment as compared with the control. The number of archaeal *mcrA* gene copies was lower ($P \leq 0.035$) with BCM and PTS from day 4 and from day 4 to 8, respectively, compared with control. No effect ($P = 0.547$) on the numbers of protozoa was observed with any treatment. However, a reduction ($P < 0.001$) of protozoa numbers was observed for all the treatments at day 4 compared to 0. The treatment x day interaction was only significant ($P < 0.001$) for the archaeal gene copy numbers.

The Roche/454 pyrosequencing analysis exhibited 21,530 and 36,183 input sequence reads of bacteria and archaea, respectively. After the removal the low-quality reads, 18,182 bacterial

and 29,404 archaeal reads were used for the analysis.

Alpha diversity based on Shannon diversity and richness of observed species measures showed similar values for bacterial diversity within the treatments, ranging from 8.1, 8.4, 8.5 and 8.7 for BCM, Control, DDS and PTS respectively on Shannon diversity and from 261, 285, 286 and 295 for BCM, Control, DDS and PTS respectively on richness of observed species. Similarly for Archaeal diversity, little variation in the observed sample diversity was observed within treatments with 4.1, 4, 4.1 and 4.1 for BCM, Control, DDS and PTS respectively on Shannon diversity and from 33, 37, 38 and 41 for BCM, Control, DDS and PTS respectively on richness of observed species.

Between treatments diversity as measured with beta diversity analysis for the comparison of the bacterial microbiome structures, found that the BCM treatment contributed the most variance to the Principal Coordinate Analysis (PCoA) (Figure 1a). Although a smaller percentage of variance was also explained in separating the PTS treated samples from the control and DDS samples. Likewise for the archaeal beta diversity, the BCM sample explained the most variance in the data (Figure 1b). However the PTS sample were also clearly separated from the control and DDS samples along the first axis of variance.

Analysis of the Bacteria family level to its specific responses to the inhibitors (based on an average of 4469 sequences per treatment) showed a substantial shift in the relative abundance of some families in fermenters treated with BCM compared to the other three treatments (Fig. 2). For BCM treatment a reduction in *Anaeroplasmataceae* (2.6%) and *Ruminococcaceae* (6.1%) compared with an average of 5.4% and 10.2%, respectively the other treatments. The BCM treatment resulted in greater abundances of *Prevotellaceae* (15.1%) and *Streptococcaceae* (3.83%) families and of an unclassified family belonging to the order *Bacteriodales* (17.1%) in comparison to the other treatments (9.1%, 1.3% and 12.3%, respectively). Treatment with PTS resulted in higher relative abundance of the

Spirochaetaceae (5.9%) family compared to the other treatments (2.63%).

Specific OTU's that were significantly associated with an increase in abundance due to treatment with BCM ($P \leq 0.05$) were classified to *Butyrivibrio* and *Prevotella* genus, while those decreasing with BCM treatment were classified as *Ruminococcus* genus (Supplementary figure 1). The PTS treatment increased ($P \leq 0.05$) the abundance of some OTUs classified as *Ruminococcus* and *Prevotella* genus. The DDS only affected ($P < 0.001$) one OTU classified as *Clostridium* genus.

Analysis of the methanogen genus level to its specific responses to the inhibitors (based on an average of 7351 sequences per treatment) (Fig. 3) revealed higher relative abundance of *Methanobrevibacter* genus in samples from fermenters treated with PTS (79.2%) and BCM (73.7%) than for control (60.6%) and DDS (48.7%) treatments. The relative abundance of archaea belonging to the *Methanosphaera* genus was higher for BCM (25.3%) than for the other three treatments (6.5%). However, the relative abundance of *Methanomicrobium* genus, from the *Methanomicrobiaceae* family was higher for control (32.9%) and DDS (41.4%) than for PTS (11.7%) and BCM (0.3%).

In relation to the archaeal community, specific OTU's that were significantly associated with an increase in abundance due to treatment with BCM and PTS ($P \leq 0.05$) were associated (Supplementary figure 2) to *Methanosphaera* and *Methanobrevibacter* genus were increased, while those decreasing with these treatments were classified as *Methanomicrobium* genus.

Discussion

In this study the compounds were provided together with the diet twice a day to mimic as close as possible the conditions *in vivo*. Thus, fluctuations in compounds concentration are expected thorough the day but no accumulation along the course of the trial. The selected dosage was made based on previous *in vitro* trials conducted using batch culture systems

(Martinez-Fernandez *et al.*, 2013). Forty ml saliva/h were infused which resulted in 960 ml/d of dilution rate. In contrast to close systems (i.e. batch culture), continuous culture do not have a steady concentration of active compound within the vessel, as it is the case of the rumen.

Ruminal fermentation

Halogenated compounds such as BCM (McAllister & Newbold, 2008), and organosulphur compounds (Patra & Yu, 2012; Mateos *et al.*, 2013) such as PTS, have shown antimethanogenic effect in ruminants. In the present work the addition of PTS and BCM decreased methane production by 48% and 98%, respectively, values that are comparable to those reported *in vitro* (Goel *et al.*, 2009; Patra & Yu, 2012; Martínez-Fernández *et al.*, 2013). On the contrary, DDS did not decrease CH₄ production from fermenters content taken on day 12. We recently reported a reduction in CH₄ production *in vitro* after 24 h treatment with DDS, with equivalent doses as used here (Martínez-Fernández *et al.*, 2013). The lack of effect of DDS on methane production using fermenter content taken on day 12 of treatment may suggest an adaptation of the microbial ecosystem over this period to the presence of this compound, which agrees with the results obtained in sheep by Klevenhusen *et al.* (2011), which suggested that some antimicrobial additives may be degraded by rumen microorganisms. The analysis of the microbial community structure in the first days of application of DDS would elucidate the adaptation mechanisms. The antimethanogenic effect of PTS observed in the present work agrees with previous results obtained *in vitro* and *in vivo* in our group (Martínez-Fernández *et al.*, 2013). The significance of the treatment by time interaction for some individual VFA molar proportions suggests that the rumen microbial community is altering the fermentation profiles either through functional changes or population shifts in the presence of the studied compounds. Total VFA concentration and

profile was only affected by BCM, in contrast to other observations (Goel *et al.*, 2009; Mitsumori *et al.*, 2012; Abecia *et al.*, 2012), which could be due to the higher dose applied here (136 μ M) as compared to Goel *et al.* (2009) and Abecia *et al.* (2012) (10 and 23 μ M, respectively). Busquet *et al.* (2005) reported no effect of DDS and garlic oil after 9 days of incubation in continuous-culture fermenters on total VFA and modifications of VFA profile with decreased acetate and increased butyrate molar proportions. In the present work acetate:propionate ratio decreased with BCM from day 4 to 12, in agreement with results obtained in previous experiments using the same compound (Mitsumori *et al.*, 2012; Abecia *et al.*, 2012). This reduction has been considered (McAllister & Newbold, 2008) a common feature for antimethanogenic compounds, as a result of a redirection of hydrogen from methane to propionic metabolic pathways, although the decrease in methane production not necessarily always result in an increase of the propionate production. In this line, some studies (Mitsumori *et al.*, 2012; Martínez-Fernández *et al.*, 2014) have reported that most of the hydrogen produced in excess in the rumen when methanogenesis is inhibited is expelled by the animal, which it is very likely to have occurred in our experiment. The potential increase in propionate would result in more energy available for the animal (Abecia *et al.*, 2012), although this effect might depend on the shifts caused in the bacterial populations as discussed below.

The treatment with PTS tended to decrease xylanase activity, while BCM and DDS did not show such effect. Xylanase degrades the linear polysaccharide beta-1,4-xylan into xylose, thus breaking down hemicellulose, one of the major components of plant cell walls in forages. The impact that this could have on plant fibre degradation in the rumen and on intakes by animals subjected to high feeding levels deserves further attention. On the other hand, only BCM treatment increased amylolytic activity on days 4 and 12. Other authors (Hristov *et al.*, 2003) reported decreased amylolytic activity by several bioactive agents using

bovine rumen fluid, in contrast to our results. These differences could be due to the different activity, chemical structure or mechanism of action of the compounds used in both studies. Janssen (2010) described that cellulolytic microbes produce more acetate and H₂, while amylolytic microbes produce less H₂ and more propionate, which explains why more CH₄ is formed, per unit of fermented matter, from forage based diets as compared to those including concentrate. That could explain the different mechanisms of action of PTS and BCM, so BCM could affect competitors of amylolytic microorganisms, thus increases amylolytic activity and produces less H₂, while PTS could affect other microbial groups as discussed in following sections.

Microbial community analyses

In agreement with the methane reduction observed with the addition of PTS and BCM, the abundance of methanogenic archaea was decreased on days 4 and 8 (and on day 12 only with BCM). Goel *et al.* (2009) reported a sharp decrease in biomass of methanogenic archaea as a result of adding BCM in batch and CCF, in agreement with our results. On the contrary Abecia *et al.* (2012) reported no changes in the concentration of methanogenic archaea in goats treated with BCM for 60 days. This disagreement could be due to the different duration of the treatments (Williams *et al.*, 2009) and the inherent differences between *in vivo* and *in vitro* conditions (Soto *et al.*, 2012; 2013). Some works reported that adding garlic compounds does not induce changes in the abundance of methanogenic archaea (Ohene-Adjei *et al.*, 2008; Kongmun *et al.*, 2011) while others reported decreased archaeal population size (Patra & Yu, 2012). The variability in concentration of active compounds in plant extracts generates confusion because the effects can be contradictory, according to the content of the active component in the extract and the dose used (Patra & Saxena, 2009). Therefore, it seems necessary either to report concentrations and active compounds in plant extracts or to use

pure products to unequivocally define activities, doses, and mechanisms of action.

The taxonomic assignment of bacteria revealed that the most abundant families were *Lachnospiraceae*, *Ruminococcaceae*, *Prevotellaceae* and unclassified *Bacteroidales* in accordance with other studies (Kong *et al.*, 2010; Zened *et al.*, 2012). The treatment with BCM increased the relative abundance of *Prevotella* and decreased those belonging to *Ruminococcus*, which is in the line of what Mitsumori *et al.* (2012) observed using the same compound. Increased abundance of *Prevotella* promoted by treatment with BCM is likely associated to the increase in branched chain fatty acids, propionate and amylolytic activity observed in that study. The abundance of *Prevotella* has been linked to hydrogen accumulation due to decreased methane production. Likewise, the effect of BCM on *Ruminococcus* is in agreement with Mitsumori *et al.* (2012) that reported a decrease in *R. albus* as a result of treating goats with BCM, due to the high sensitivity to high partial pressure of hydrogen. This decrease might be compensated by greater abundances of other fibrolytic bacteria such as *F. succinogenes* that does not produce H₂ and is not susceptible to H₂ accumulation. This agrees with Kittelmann *et al.* (2014), who reported that two different bacterial community types are linked with the low-methane emission trait in sheep, hypothesizing that lower CH₄ yields are the result of bacterial communities that ferment ingested feed to relatively less hydrogen, which results in less CH₄ being formed. Furthermore, BCM interferes with the cobalamin-dependent pathway to reduce CO₂ to CH₄ and thus decreasing methanogenesis. Some cellulolytic bacteria have been shown to be dependent on vitamin B12 (Scott & Dehority, 1965), which could explain the shift in the distribution of some bacterial groups observed here and the reduction in the abundance of *Ruminococcus* when treating with BCM. With regards to PTS, although a decrease in methane production was observed, it did not induce detectable changes in the bacterial taxonomic distribution, which could be explained by the fact that the reduction in methane

was not as dramatic as for BCM (48 and 94 %, respectively for PTS and BCM) and theoretically no major shift in metabolic H₂ transfer occurred. Indeed, this is confirmed by the PCoA plot in which a distinct group including BCM samples was recognized and separated from the rest.

The dominant archaea belonged to the orders *Methanobacteriales* and *Methanomicrobiales*, in accordance with previous works (Janssen & Kirs, 2008; Zhou *et al.*, 2009; GU *et al.*, 2011). Both BCM and PTS increased the relative abundance of *Methanobrevibacter* and decreased that of *Methanomicrobium* compared with control and DDS. The BCM treatment also increased archaea from *Methanosphaera*. These results are in agreement with previous results obtained in goats treated with BCM for two months (Abecia *et al.*, 2014). At least 60 genes are involved in methanogenesis and hydrogen transfer. The first five steps of the pathway result in the sequential reduction of CO₂ by electrons sourced from H₂ to form N⁵-methyl-H₄MPT (Thauer *et al.*, 1993). The methyl group is then transferred to coenzyme M via the action of methyl-H₄MPT:CoM-methyltransferase which is encoded by the *mtr* gene cluster and this is the step inhibited by BCM. This is a multi-subunit enzyme encoded by more than 10 different genes (Attwood & McSweeney, 2008) that include the methyl CoM reductase cluster (Mcr B, D, C, G, A) and the CoM methyltransferase cluster (Mtr E, D, C, B, A, G, H, X) and a set of genes whose function is currently not known. Within the hydrogenotrophic methanogens, a further 10 genes are conserved (Gao & Gupta, 2007) and include a cluster of genes that encode subunits of [Ni-Fe] hydrogenases (Eha B, C, D, E, F, G) that catalyse the reversible reduction of protons to molecular hydrogen. Although some of the genes are known to be conserved across methanogens, some differ between families (Attwood & McSweeney, 2008). Since our results are consistent with other work using BCM (Abecia *et al.*, 2014), we hypothesize that the different sensitivity of *Methanobacteriales* and *Methanomicrobiales* is a result of genes differently expressed that make some species more

suitable to cope with the new environment.

Our results support previous observations by Ohene-Adjei *et al.* (2008) who reported changes in the archaeal banding profile by DGGE as a result of treating sheep with garlic oil. Based on the taxonomy derived from pyrosequencing we hypothesize that the extent in which groups are shifted is directly associated to the extent of methane inhibition. However, the different mechanisms of action of BCM and PTS could also explain the different impact observed. Bromochloromethane directly reacts with reduced vitamin B12 and results in the inhibition of cobamide-dependent methyl group transfer in methanogenesis (Wood *et al.*, 1968), while the antimicrobial effect of thiosulfates (Focke *et al.*, 1990; Ruiz *et al.*, 2010) is associated with chemical reaction with thiol groups of various enzymes such as the acetyl-CoA-forming system. Some authors (Busquet *et al.*, 2005; Benchaar & Greathead, 2011) reported the relationship of antimethanogenic effect of organosulphur and the inhibition of HMG-CoA reductase, which play an important role in the synthesis of isoprenoid ethers, the main component of archaeal cell membranes. Therefore, the sensitivity of key archaeal groups to these inhibitory compounds may be explained by the different mechanisms of action exhibited, as discussed above, which would need to be further tested using deep metagenomic and metatranscriptomic sequencing transcription studies combined with pure culture *in vitro* incubations. Kang *et al.*, (2013) used a combined RNA and DNA-derived analysis and concluded that less abundant but highly active methanogens may make a greater contribution to total methane formation than their abundance may suggest. Shi *et al.* (2014) have recently observed that in despite of minor changes in the abundance of different methanogens groups in the rumen of sheep that are consistently high or low CH₄ yield phenotypes, the transcription of methanogenesis pathway genes was substantially different. This differential transcription pattern needs to be assessed in the future when effective anti-methanogenesis compounds are applied.

In conclusion, the inhibition of methane production in the rumen by BCM (94%) and PTS (48%) is associated with a shift in the archaeal biodiversity that involves an increase in *Methanospaera* and a decrease in *Methanomicrobium*. In the case of BCM the effect causes changes in the bacterial population that are clearly reflected in the fermentation profile. The potential negative impact of PTS on fibre degradation deserves further studies. The treatment with DDS over 12 days does not confirm the antimethanogenic effect previously reported.

Acknowledgement

This research was supported by the Science and Innovation Spanish Ministry (Project AGL2008-04707-C02-01). We gratefully acknowledge T. García, E. Jiménez and P. Rufino for their technical assistance and DMC Research Center S.L. for providing PTS. G. Martínez gratefully acknowledges the FPI grant from the Spanish Science Ministry.

References

- Abecia L, Toral P, Martín-García A, Martínez G, Tomkins N, Molina-Alcaide E, Newbold C & Yáñez-Ruiz D (2012) Effect of bromochloromethane on methane emission, rumen fermentation pattern, milk yield, and fatty acid profile in lactating dairy goats. *Journal of Dairy Science* **95**: 2027-2036.
- Abecia L, Waddams KE, Martínez-Fernandez G, Martín-García AI, Ramos-Morales E, Newbold CJ & Yáñez-Ruiz DR (2014) An antimethanogenic nutritional intervention in early life of ruminants modifies ruminal colonization by Archaea. *Archaea*, 2014.
- Association of Official Analytical Chemists (AOAC) (2005) Official Methods of Analysis 18th ed. AOAC, Gaithersburg, MD.

- 493 Attwood G & McSweeney C (2008) Methanogen genomics to discover targets for methane
494 mitigation technologies and options for alternative H₂ utilisation in the rumen.
495 *Australian Journal of Experimental Agriculture*, **48**: 28–37.
- 496 Benchaar C & Greathead H (2011) Essential oils and opportunities to mitigate enteric
497 methane emissions from ruminants. *Animal Feed Science and Technology* **166–167**:
498 338–355.
- 499 Bragg L, Stone G, Imelfort M, Hugenholtz P & Tyson GW (2012) Fast, accurate error-
500 correction of amplicon pyrosequences using Acacia. *Nature Methods* **9**: 425–426.
- 501 Busquet M, Calsamiglia S, Ferret A, Carro M & Kamel C (2005) Effect of garlic oil and four
502 of its compounds on rumen microbial fermentation. *Journal of Dairy Science* **88**:
503 4393–4404.
- 504 Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL & Knight R (2010a)
505 PyNAST: a flexible tool for aligning sequences to a template alignment.
506 *Bioinformatics* **26**: 266–267.
- 507 Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N,
508 Pena AG, Goodrich JK & Gordon JI (2010b) QIIME allows analysis of high-
509 throughput community sequencing data. *Nature methods* **7**: 335–336.
- 510 Denman SE, Tomkins NW & McSweeney CS (2007) Quantification and diversity analysis of
511 ruminal methanogenic populations in response to the antimethanogenic compound
512 bromochloromethane. *FEMS Microbiology. Ecology* **62**: 313–322.
- 513 DeSantis JrT, Hugenholtz P, Keller K, Brodie E, Larsen N, Piceno Y, Phan R & Andersen G
514 (2006) NAST: a multiple sequence alignment server for comparative analysis of 16S
515 rRNA genes. *Nucleic Acids Research* **34**: W394–W399.
- 516 Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST.
517 *Bioinformatics* **26**: 2460–2461.

- 518 European Directive 86/609. Commission recommendations of 18 June 2007 on guidelines for
 519 the accommodation and care of animals used for experimental and other scientific
 520 purposes. Annex II to European Council Directive 86/609. The Commission of the
 521 European Communities Publishing, Brussels, Belgium.
- 522 Focke M, Feld A & Lichtenthaler HK (1990) Allicin, a naturally occurring antibiotic from
 523 garlic, specifically inhibits acetyl-CoA synthetase. *FEBS letters* **261**: 106-108.
- 524 Galand PE, Casamayor EO, Kirchman DL, Potvin M & Lovejoy C (2009) Unique archaeal
 525 assemblages in the Arctic Ocean unveiled by massively parallel tag sequencing. *The*
 526 *ISME journal* **3**: 860-869.
- 527 Gao B & Gupta RS (2007) Phylogenomic analysis of proteins that are distinctive of Archaea
 528 and its main subgroups and the origin of methanogenesis. *BMC genomics*, **8**(1), 86.
- 529 Giraldo L, Tejido M, Ranilla M & Carro M (2008) Effects of exogenous fibrolytic enzymes
 530 on in vitro ruminal fermentation of substrates with different forage: concentrate ratios.
 531 *Animal Feed Science and Technology* **141**:306-325.
- 532 Goel G, Makkar HPS & Becker K (2009) Inhibition of methanogens by
 533 bromochloromethane: effects on microbial communities and rumen fermentation
 534 using batch and continuous fermentations. *British Journal of Nutrition* **101**: 1484-
 535 1492.
- 536 Gu MJ, Alam M, Kim SH, Jeon CO, Chang MB, Oh YK, Lee SC & Lee SS (2011) Analysis
 537 of methanogenic archaeal communities of rumen fluid and rumen particles from
 538 Korean black goats. *Animal Science Journal* **82**:663-672.
- 539 Hart KJ, Yáñez-Ruiz DR, Duval SM, McEwan NR & Newbold CJ (2008) Plant extracts to
 540 manipulate rumen fermentation. *Animal Feed Science and Technology* **147**: 8-35.
- 541 Isac M, García M, Aguilera J & Molina AE (1994) A comparative study of nutrient
 542 digestibility, kinetics of digestion and passage and rumen fermentation pattern in

- 543 goats and sheep offered medium quality forages at the maintenance level of feeding.
 544 *Archiv für Tierernährung* **46**: 37.
- 545 Janssen PH & Kirs M (2008) Structure of the archaeal community of the rumen. *Applied and*
 546 *environmental microbiology* **74**: 3619-3625.
- 547 Janssen PH (2010) Influence of hydrogen on rumen methane formation and fermentation
 548 balances through microbial growth kinetics and fermentation thermodynamics.
 549 *Animal Feed Science and Technology* **160**: 1-22.
- 550 Kang SH, Evans P, Morrison M & McSweeney C (2013) Identification of metabolically
 551 active proteobacterial and archaeal communities in the rumen by DNA- and RNA-
 552 derived 16S rRNA gene. *Journal of Applied Microbiology* **115**: 644– 653.
- 553 Kittelmann S, Pinares-Patiño C, Seedorf H, Kirk MR, Ganesh S., McEwan J.C & Janssen
 554 P.H (2014) Two different bacterial community types are linked with the low-methane
 555 emission trait in sheep. *PLOS One*. **9**: e103171
- 556 Klevenhusen F, Zeitz JO, Duval S, Kreuzer M & Soliva CR (2011) Garlic oil and its
 557 principal component diallyl disulfide fail to mitigate methane, but improve
 558 digestibility in sheep. *Animal Feed Science and Technology* **166–167**: 356-363.
- 559 Kong Y, Teather R & Forster R (2010) Composition, spatial distribution, and diversity of the
 560 bacterial communities in the rumen of cows fed different forages. *FEMS*
 561 *microbiology ecology* **74**: 612-622.
- 562 Kongmun P, Wanapat M, Pakdee P, Navanukraw C & Yu Z (2011) Manipulation of rumen
 563 fermentation and ecology of swamp buffalo by coconut oil and garlic powder
 564 supplementation. *Livestock Science* **135**:84-92.
- 565 Li W & Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of
 566 protein or nucleotide sequences. *Bioinformatics* **22**: 1658-1659.

- 567 Liu Z, Lozupone C, Hamady M, Bushman FD & Knight R (2007) Short pyrosequencing
568 reads suffice for accurate microbial community analysis. *Nucleic Acids Research* **35**:
569 e120.
- 570 Lozupone C & Knight R (2005) UniFrac: a new phylogenetic method for comparing
571 microbial communities. *Applied and Environmental Microbiology* **71**: 8228-8235.
- 572 Maeda H, Fujimoto C, Haruki Y, Maeda T, Kokeguchi S, Petelin M, Arai H, Tanimoto I,
573 Nishimura F & Takashiba S (2003) Quantitative real-time PCR using TaqMan and
574 SYBR Green for *Actinobacillus actinomycetemcomitans*, *Porphyromonas gingivalis*,
575 *Prevotella intermedia*, *tetQ* gene and total bacteria. *FEMS Immunology & Medical*
576 *Microbiology* **39**(1): 81-86.
- 577 Martínez-Fernández G, Abecia L, Martín-García AI, Ramos-Morales E, Hervás G, Molina-
578 Alcaide E & Yáñez-Ruiz DR (2013) In vitro–in vivo study on the effects of plant
579 compounds on rumen fermentation, microbial abundances and methane emissions in
580 goats. *Animal* **7**: 1925-1934.
- 581 Martínez-Fernández G, Tomkins NW, Padmanabha J, Denman SE & McSweeney CS (2014)
582 Rumen Methane Reduction Mediated by Chloroform Increases Expelled Hydrogen in
583 Cattle. *Proceedings of the 30th Biennial Conference of the Australian Society for*
584 *Animal Production* **30**: 49. ISSN 0728-5965
- 585 Mateos I, Ranilla M, Tejido M, Saro C, Kamel C & Carro MD (2013) The influence of diet
586 type (dairy versus intensive fattening) on the effectiveness of garlic oil and
587 cinnamaldehyde to manipulate in vitro ruminal fermentation and methane production.
588 *Animal Production Science* **53**: 299-307.
- 589 May C, Payne AL, Stewart PL & Edgar JA (1995) A delivery system for agents. AU
590 International Patent PCT/AU95/00733. Australian Industrial Property

- 591 Organisation, Canberra, ACT, Australia.
- 592 McAllister T & Newbold CJ (2008) Redirecting rumen fermentation to reduce
 593 methanogenesis. *Australian Journal of Experimental Agriculture* **48**: 7-13.
- 594 Mitsumori M, Shinkai T, Takenaka A, Enishi O, Higuchi K, Kobayashi Y, Nonaka I,
 595 Asanuma N, Denman SE & McSweeney CS (2012) Responses in digestion, rumen
 596 fermentation and microbial populations to inhibition of methane formation by a
 597 halogenated methane analogue. *British Journal of Nutrition* **108**: 482-491.
- 598 Morgavi D, Forano E, Martin C & Newbold CJ (2010) Microbial ecosystem and
 599 methanogenesis in ruminants. *Animal* **4**: 1024-1036.
- 600 Muetzel S, Lawrence P, Hoffmann EM & Becker K (2009) Evaluation of a stratified
 601 continuous rumen incubation system. *Animal Feed Science and Technology* **151**:32-
 602 43.
- 603 Ohene-Adjei S, Chaves A, McAllister T, Benchaar C, Teather R & Forster R (2008)
 604 Evidence of increased diversity of methanogenic archaea with plant extract
 605 supplementation. *Microbial Ecology* **56**:234-242.
- 606 Patra AK & Saxena J (2009) Dietary phytochemicals as rumen modifiers: a review of the
 607 effects on microbial populations. *Antonie van Leeuwenhoek* **96**: 363-375.
- 608 Patra AK & Yu Z (2012) Effects of Essential Oils on Methane Production and Fermentation
 609 by, and Abundance and Diversity of, Rumen Microbial Populations. *Applied and*
 610 *Environmental Microbiology* **78**: 4271-4280.
- 611 Royal Decree No. 1201/2005 of 10 October on the protection of animals used for
 612 experimentation and other scientific purposes. Boletín del Estado, pp. 34367-34391.
 613 Spanish Government Publishing, Madrid, Spain.

- 614 Ruiz R, García M, Lara A & Rubio L (2010) Garlic derivatives (PTS and PTS-O) differently
 615 affect the ecology of swine faecal microbiota in vitro. *Veterinary Microbiology* **144**:
 616 110-117.
- 617 Scott HW & Dehority BA (1965) Vitamin Requirements of Several Cellulolytic Rumen
 618 Bacteria. *Journal of Microbiology* **18**: 1169-1175.
- 619 Shi W, Moon CD, Leahy SC, Kang D, Froula J, Kittelmann S, ... & Rubin EM (2014)
 620 Methane yield phenotypes linked to differential gene expression in the sheep rumen
 621 microbiome. *Genome Research* **24**: 1517-25
- 622 Soliva CR, Amelchanka SL, Duval SM & Kreuzer M (2011) Ruminal methane inhibition
 623 potential of various pure compounds in comparison with garlic oil as determined with
 624 a rumen simulation technique (Rusitec). *British Journal of Nutrition* **106**: 114-122.
- 625 Soto E, Yáñez-Ruiz D, Cantalapiedra-Hijar G, Vivas A & Molina-Alcaide E (2012) Changes
 626 in ruminal microbiota due to rumen content processing and incubation in single-flow
 627 continuous-culture fermenters. *Animal Production Science* **52**: 813-822.
- 628 Soto E, Molina-Alcaide E, Khelil H & Yáñez-Ruiz D (2013) Ruminal microbiota developing
 629 in different in vitro simulation systems inoculated with goats' rumen liquor. *Animal*
 630 *Feed Science and Technology* **185**: 9-18.
- 631 Sylvester JT, Karnati SK, Yu Z, Morrison M & Firkins JL (2004) Development of an assay to
 632 quantify rumen ciliate protozoal biomass in cows using real-time PCR. *The Journal of*
 633 *Nutrition* **134**:3378–3384.
- 634 Thauer RK, Hedderich R & Fischer R (1993) Reactions and enzymes involved in
 635 methanogenesis from CO₂ and H₂. *Methanogenesis: Ecology, Physiology,*
 636 *Biochemistry and Genetics* (Ferry JG, ed), pp. 209–252, Chapman & Hall, New York,
 637 NY, USA

- 638 Theodorou MK, Williams BA, Dhanoa MS, McAllan AB & France J. (1994) A simple gas
639 production method using a pressure transducer to determine the fermentation kinetics
640 of ruminant feeds. *Animal Feed Science and Technology* **48**:185-197.
- 641 Van Soest PJ, Robertson J & Lewis B (1991) Methods for dietary fiber, neutral detergent
642 fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy*
643 *Science* **74**:3583-3597.
- 644 Zened A, Combes S, Cauquil L, Mariette J, Klopp C, Bouchez O, Troegeler-Meynadier A &
645 Enjalbert F (2012) Microbial ecology of the rumen evaluated by 454 GS FLX
646 pyrosequencing is affected by starch and oil supplementation of diets. *FEMS*
647 *Microbiology Ecology* **83**: 504-514
- 648 Zhou M, Hernandez-Sanabria E & Guan LL (2009) Assessment of the microbial ecology of
649 ruminal methanogens in cattle with different feed efficiencies. *Applied and*
650 *Environmental Microbiology* **75**: 6524-6533.
- 651

651 Table 1. Chemical composition of alfalfa hay and concentrate (g kg^{-1} dry matter) and
 652 ingredients (g kg^{-1}) of concentrate.

Item	Alfalfa hay	Concentrate
DM (g kg^{-1} fresh matter)	907	915
OM	875	884
NDF	513	245
ADF	330	118
ADL	99.2	36.3
CP	203	168
Ether Extract	8.1	15.3
GE (MJ Kg^{-1} DM)	18.4	19.5
Ingredients		
Barley		174
Faba beans		233
Maize		174
Sunflower meal		233
Maize gluten meal		116
Rumen-inert fat		70

653

654

655

Table 2. Effects of additives on VFA concentration (mM), profiles (mol 100⁻¹ mol) and pH on days 0, 4, 8 and 12 of incubation in continuous-culture fermenters and on CH₄ production (μmol) after 24 hours of incubation in batch culture inoculated with fermenters content after 12 days of incubation.

Item	Day	Treatment				SEM ¹	P-value ²		
		Control	BCM	DDS	PTS		T	d	Txd
Total VFA	0	30.6 ^B	31.6 ^B	28.8 ^B	28.7 ^B	11.4	0.015	<0.001	0.013
	4	76.2 ^{aA}	62.9 ^{bA}	74.9 ^{aA}	73.1 ^{aA}				
	8	73.9 ^{aA}	60.2 ^{bA}	74.82 ^{aA}	72.4 ^{aA}				
	12	77.4 ^{aA}	65.7 ^{bA}	78.1 ^{aA}	78.9 ^{aA}				
Acetate	0	69.0 ^A	68.4 ^A	68.2 ^A	68.4 ^A	1.8	<0.001	<0.001	<0.001
	4	63.7 ^{aB}	50.0 ^{cB}	61.4 ^{bB}	62.8 ^{abB}				
	8	64.9 ^{aB}	49.4 ^{cB}	62.6 ^{bB}	62.6 ^{bB}				
	12	64.6 ^{aB}	49.6 ^{bB}	61.9 ^{aB}	62.1 ^{aB}				
Propionate	0	13.8 ^B	14.1 ^C	14.1 ^B	13.9 ^B	1.5	<0.001	<0.001	<0.001
	4	16.0 ^{bA}	25.8 ^{aA}	17.6 ^{bA}	17.1 ^{bA}				
	8	17.0 ^{cA}	24.1 ^{aA}	17.5 ^{bA}	17.8 ^{bA}				
	12	17.8 ^{bA}	23.4 ^{aB}	18.6 ^{bA}	18.0 ^{bA}				
Isobutyrate	0	1.88 ^A	1.85 ^A	1.80 ^A	1.83 ^A	0.16	0.163	0.013	0.118
	4	1.48 ^B	1.23 ^B	1.38 ^B	1.43 ^B				
	8	1.33 ^B	1.25 ^B	1.35 ^B	1.23 ^B				
	12	1.30 ^B	1.22 ^B	1.35 ^B	1.35 ^B				
Butyrate	0	11.2	11.4 ^B	12.1 ^B	11.9	0.7	0.002	<0.001	<0.001
	4	13.8 ^b	16.3 ^{aA}	14.5 ^{bA}	13.9 ^b				
	8	12.2 ^b	15.4 ^{aA}	13.3 ^{bAB}	13.8 ^{ab}				
	12	11.7 ^c	15.6 ^{aA}	12.9 ^{bcAB}	13.7 ^b				
Isovalerate	0	2.93	3.03	2.70	2.83	0.39	<0.001	0.405	<0.001
	4	2.78	2.73	2.75	2.50				
	8	2.38 ^b	5.38 ^a	2.68 ^b	2.20 ^b				
	12	2.38 ^b	5.90 ^a	2.58 ^b	2.33 ^b				
Valerate	0	1.28 ^B	1.28 ^B	1.25 ^C	1.20 ^B	0.32	<0.001	<0.001	<0.001
	4	2.20 ^{bA}	4.00 ^{aA}	2.38 ^{bB}	2.35 ^{bA}				
	8	2.20 ^{cA}	4.43 ^{aA}	2.50 ^{bAB}	2.38 ^{bcA}				
	12	2.33 ^{bA}	4.38 ^{aA}	2.68 ^{bA}	2.55 ^{bA}				
Acetate:Propionate	0	5.02 ^A	4.88 ^A	4.90 ^A	4.97 ^A	0.46	<0.001	<0.001	<0.001
	4	3.98 ^{aB}	1.94 ^{dB}	3.48 ^{cB}	3.67 ^{bB}				
	8	3.85 ^{aB}	2.06 ^{bB}	3.57 ^{aB}	3.52 ^{aB}				
	12	3.67 ^{aB}	2.13 ^{cB}	3.34 ^{bB}	3.46 ^{abB}				
pH	0	6.55 ^A	6.58	6.55 ^A	6.58	0.22	0.138	0.033	0.668
	4	6.39 ^{AB}	6.45	6.43 ^{AB}	6.39				
	8	6.31 ^B	6.44	6.41 ^{AB}	6.40				
	12	6.36 ^{AB}	6.44	6.38 ^B	6.40				
Methane	13	249 ^a	14 ^c	248 ^a	129 ^b	13	<0.001	n.d.	n.d.

660 Treatment: control (without additive), DDS (diallyl disulfide), PTS (propyl propane
661 thiosulfinate) and BCM (Bromochloromethane).

662 ¹SEM: Standard error of the mean.

663 ² T: treatment effect; d: day effect; Txd: Treatment x day interaction.

664 ^{a-c} within a row treatment means without a common superscript differ, $P < 0.05$.

665 ^{A-B} within a column treatment without a common superscript differ, $P < 0.05$.

666

667

Table 3. Effect of the additives on xylanase and amylase activities in CCF content sampled on day 4 and 12 after inoculation.

Item	Day	Treatment				SEM ¹	P-value ²		
		Control	BCM	DDS	PTS		T	d	Txd
Xylanase	4	5.72	6.52	5.65	5.14	1.58	0.051	0.458	0.186
	12	6.91	7.10	6.08	5.22				
Amylase	4	1.03 ^b	1.33 ^a	1.02 ^b	1.04 ^b	0.23	0.022	0.571	0.867
	12	1.10 ^b	1.23 ^a	1.06 ^b	1.08 ^b				

669

670 Treatment: control (without additive), DDS (diallyl disulfide), PTS (propyl propane
671 thiosulfinate) and BCM (Bromochloromethane).

672 ¹SEM: Standard error of the mean.

673 ² T: treatment effect; d: day effect; Txd: Treatment x day interaction. ^{a-c} within a row

674 treatment means without a common superscript differ, $P < 0.05$.

675 Amylase activity is expressed as micromoles of glucose released from soluble starch by 1 mL
676 of ruminal fluid in 1 min at 39°C and pH=6.5. Xylanase activity is expressed as micromoles
677 of xylose liberated from oat beachwood xylan by 1 mL of ruminal fluid in 1 min at 39°C and
678 pH=6.5.

679

Table 4. Effects of the additives on the concentration (log copy gene numbers g⁻¹ fresh matter) of total bacteria (16S rRNA), protozoa (18S rRNA) and methanogenic archaea (mcrA gene) in fermenters after 0, 4, 8 and 12 days of incubation.

Item	Day	Treatment				SEM ¹	P-value ²		
		Control	BCM	DDS	PTS		T	d	Txd
Methanogens	0	8.72	8.63 ^A	8.69 ^{AB}	8.68 ^A	0.10	<0.001	0.027	<0.001
	4	8.62 ^a	7.91 ^{bB}	8.17 ^{abB}	7.96 ^{abAB}				
	8	8.69 ^a	7.19 ^{cC}	8.37 ^{bAB}	8.26 ^{bB}				
	12	8.81 ^a	6.90 ^{bC}	8.71 ^{aA}	8.26 ^{aAB}				
Protozoa	0	8.11 ^A	7.96 ^A	8.14 ^A	8.05 ^A	0.51	0.551	<0.001	0.033
	4	7.23 ^B	7.13 ^B	7.28 ^B	6.96 ^B				
	8	6.93 ^B	7.02 ^B	6.48 ^C	6.74 ^B				
	12	6.79 ^B	6.99 ^B	6.63 ^{BC}	6.92 ^B				
Bacteria	0	9.86 ^A	9.82 ^A	9.88 ^A	9.79 ^A	0.09	0.010	0.002	0.099
	4	9.49 ^{bB}	9.76 ^{aA}	9.58 ^{abB}	9.54 ^{bABC}				
	8	9.50 ^B	9.56 ^B	9.45 ^B	9.44 ^C				
	12	9.51 ^B	9.74 ^A	9.53 ^B	9.68 ^{AB}				

Treatment: control (without additive), DDS (diallyl disulfide), PTS (propyl propane thiosulfinate) and BCM (Bromochloromethane).

¹SEM: Standard error of the mean.

² T: treatment effect; d: day effect; Txt: Treatment x day interaction.

^{a-c} within a row treatment means without a common superscript differ, $P < 0.05$.

^{A-B} within a column treatment without a common superscript differ, $P < 0.05$.





Figure 1. Principal coordinate analysis showing the relationships of weighted bacterial (a) and archaeal (b) communities with jackknife support of the fermenters content treated with BCM (Red), PTS (Green), DDS (Orange) and without treatment (Blue) after 12 days of incubation. Size of spots represent robustness of principal coordinate analysis based on jackknife for 1000 subset resamplings.

697



698

699

700

701

702

703

704

Figure 2. Bacterial taxonomic composition of the fermenters content without treatment (Control), or treated with BCM, DDS and PTS after 12 days of incubation at family level. Sequences were classified using BLAST with a 97% similarity level. Further information regarding the complete family level taxonomic classification is available in supplementary table 1.




Figure 3. Archaeal taxonomic composition of the fermenters content without treatment (Control), or treated with BCM, DDS and PTS after 12 days of incubation. Sequences were classified using BLAST with a 97% similarity level. Further information regarding the complete family level taxonomic classification is available in supplementary table 2. * Conflict of taxonomy for GreenGenes database. They were most associated to *Methanomicrobium mobile*.